

## Eating Attitudes Test (EAT-26)<sup>®</sup>

Instructions: This is a screening measure to help you determine whether you might have an eating disorder that needs professional attention. This screening measure is not designed to make a diagnosis of an eating disorder or take the place of a professional consultation. Please fill out the below form as accurately, honestly and completely as possible. There are no right or wrong answers. All of your responses are confidential.

### Part A: Complete the following questions:

1) Birth Date	Month: <input style="width: 40px;" type="text"/>	Day: <input style="width: 40px;" type="text"/>	Year: <input style="width: 60px;" type="text"/>	2) Gender:	Male <input type="checkbox"/>	Female <input type="checkbox"/>
3) Height	Feet : <input style="width: 40px;" type="text"/>	Inches: <input style="width: 40px;" type="text"/>			<input type="checkbox"/>	<input type="checkbox"/>
4) Current Weight (lbs.):	5) Highest Weight (excluding pregnancy): <input style="width: 100px;" type="text"/>					
6) Lowest Adult Weight:	7: Ideal Weight: <input style="width: 100px;" type="text"/>					

### Part B: Check a response for each of the following statements:

	Always	Usually	Often	Some times	Rarely	Never
1. Am terrified about being overweight.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2. Avoid eating when I am hungry.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3. Find myself preoccupied with food.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4. Have gone on eating binges where I feel that I may not be able to stop.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
5. Cut my food into small pieces.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
6. Aware of the calorie content of foods that I eat.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
7. Particularly avoid food with a high carbohydrate content (i.e. bread, rice, potatoes, etc.)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
8. Feel that others would prefer if I ate more.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
9. Vomit after I have eaten.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
10. Feel extremely guilty after eating.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
11. Am preoccupied with a desire to be thinner.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
12. Think about burning up calories when I exercise.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
13. Other people think that I am too thin.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
14. Am preoccupied with the thought of having fat on my body.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
15. Take longer than others to eat my meals.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
16. Avoid foods with sugar in them.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
17. Eat diet foods.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
18. Feel that food controls my life.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
19. Display self-control around food.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
20. Feel that others pressure me to eat.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
21. Give too much time and thought to food.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
22. Feel uncomfortable after eating sweets.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
23. Engage in dieting behavior.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
24. Like my stomach to be empty.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
25. Have the impulse to vomit after meals.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
26. Enjoy trying new rich foods.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

### Part C: Behavioral Questions:

#### In the past 6 months have you:

		Never	Once a month or less	2-3 times a month	Once a week	2-6 times a week	Once a day or more
A	Gone on eating binges where you feel that you may not be able to stop? *	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
B	Ever made yourself sick (vomited) to control your weight or shape?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
C	Ever used laxatives, diet pills or diuretics (water pills) to control your weight or shape?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
D	Exercised more than 60 minutes a day to lose or to control your weight?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
E	Lost 20 pounds or more in the past 6 months	Yes <input type="checkbox"/>		No <input type="checkbox"/>			

\* Defined as eating much more than most people would under the same circumstances and feeling that eating is out of control

## Eating Attitudes Test (EAT-26)<sup>©</sup> Item Scoring

Instructions: This is a screening measure to help you determine whether you might have an eating disorder that needs professional attention. This screening measure is not designed to make a diagnosis of an eating disorder or take the place of a professional consultation. Please fill out the below form as accurately, honestly and completely as possible. There are no right or wrong answers. All of your responses are confidential.

### Part A: Complete the following questions:

1) Birth Date	Month: <input style="width: 50px;" type="text"/>	Day: <input style="width: 50px;" type="text"/>	Year: <input style="width: 50px;" type="text"/>	2) Gender:	Male <input type="checkbox"/>	Female <input type="checkbox"/>
3) Height	Feet : <input style="width: 50px;" type="text"/>	Inches: <input style="width: 50px;" type="text"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4) Current Weight (lbs.):	<input style="width: 100px;" type="text"/>	5) Highest Weight (excluding pregnancy):	<input style="width: 100px;" type="text"/>			
6) Lowest Adult Weight:	<input style="width: 100px;" type="text"/>	7: Ideal Weight:	<input style="width: 100px;" type="text"/>			

Part B: Check a response for each of the following statements:	Always	Usually	Often	Some times	Rarely	Never
1. Am terrified about being overweight.	3	2	1	0	0	0
2. Avoid eating when I am hungry.	3	2	1	0	0	0
3. Find myself preoccupied with food.	3	2	1	0	0	0
4. Have gone on eating binges where I feel that I may not be able to stop.	3	2	1	0	0	0
5. Cut my food into small pieces.	3	2	1	0	0	0
6. Aware of the calorie content of foods that I eat.	3	2	1	0	0	0
7. Particularly avoid food with a high carbohydrate content (i.e. bread, rice, potatoes, etc.)	3	2	1	0	0	0
8. Feel that others would prefer if I ate more.	3	2	1	0	0	0
9. Vomit after I have eaten.	3	2	1	0	0	0
10. Feel extremely guilty after eating.	3	2	1	0	0	0
11. Am preoccupied with a desire to be thinner.	3	2	1	0	0	0
12. Think about burning up calories when I exercise.	3	2	1	0	0	0
13. Other people think that I am too thin.	3	2	1	0	0	0
14. Am preoccupied with the thought of having fat on my body.	3	2	1	0	0	0
15. Take longer than others to eat my meals.	3	2	1	0	0	0
16. Avoid foods with sugar in them.	3	2	1	0	0	0
17. Eat diet foods.	3	2	1	0	0	0
18. Feel that food controls my life.	3	2	1	0	0	0
19. Display self-control around food.	3	2	1	0	0	0
20. Feel that others pressure me to eat.	3	2	1	0	0	0
21. Give too much time and thought to food.	3	2	1	0	0	0
22. Feel uncomfortable after eating sweets.	3	2	1	0	0	0
23. Engage in dieting behavior.	3	2	1	0	0	0
24. Like my stomach to be empty.	3	2	1	0	0	0
25. Have the impulse to vomit after meals.	3	2	1	0	0	0
26. Enjoy trying new rich foods.	0	0	0	1	2	3

Part C: Behavioral Questions: In the past 6 months have you:		Never	Once a month or less	2-3 times a month	Once a week	2-6 times a week	Once a day or more
A	Gone on eating binges where you feel that you may not be able to stop? *			√	√	√	√
B	Ever made yourself sick (vomited) to control your weight or shape?		√	√	√	√	√
C	Ever used laxatives, diet pills or diuretics (water pills) to control your weight or shape?		√	√	√	√	√
D	Exercised more than 60 minutes a day to lose or to control your weight?						√
E	Lost 20 pounds or more in the past 6 months	Yes	√	No			

\* Defined as eating much more than most people would under the same circumstances and feeling that eating is out of control

© Copyright: EAT-26: (Garner et al. 1982, *Psychological Medicine*, 12, 871-878); adapted by D. Garner with permission.

# Eating Attitudes Test<sup>®</sup> (EAT-26): Scoring and Interpretation

David M. Garner, Ph. D.

The Eating Attitudes Test (EAT-26) is probably the most widely used standardized measure of symptoms and concerns characteristic of eating disorders (Garner & Garfinkel, 1979; Garner, Olmsted, Bohr, & Garfinkel, 1982). The original EAT appeared as a Current Contents Citation Classic in 1993 (Garner, 1993). The 26-item version (Garner et al., 1989) is highly reliable and valid (Garner, Olmsted, Bohr, & Garfinkel, 1982; Lee et al., 2002; Mintz & O'Halloran, 2000). The EAT-26 alone does not yield a specific diagnosis of an eating disorder (neither the EAT-26, nor any other screening instrument, has been established as highly efficient as the sole means for identifying eating disorders).

Nevertheless, many studies have used the EAT-26 as an economical first step in a two-stage screening process. According to this methodology, individuals who score 20 or more on the test should be interviewed by a qualified professional to determine if they meet the diagnostic criteria for an eating disorder (Dotti & Lazzari, 1998; Patton, Johnson-Sabine, Wood, Mann, & Wakeling, 1990). If you have a low score on the EAT-26 (below 20), you still could have a serious eating problem, so do not let the results deter you from seeking help. The EAT-26 can be used in group or individual settings and is designed to be self-administered or be administered by health professionals, school counselors, coaches, camp counselors, and others with interest in gathering information to determine if an individual should be referred to a specialist for evaluation for an eating disorder.

The EAT-26 has been particularly useful a screening tool to assess "eating disorder risk" in high school, college and other special risk samples such as athletes (Garner, Rosen and Barry, 1998). Screening for eating disorders is based on the assumption that early identification of an eating disorder can lead to earlier treatment, thereby reducing serious physical and psychological complications or even death.

The EAT-26 items form three subscales: 1) Dieting, 2) Bulimia and Food Preoccupation, and 3) Oral Control. The subscale scores are computed by summing all items assigned to that particular scale:

*Dieting scale items:* 1, 6, 7, 10, 11, 12, 14, 16, 17, 22, 23, 24, 26.

*Bulimia & Food Preoccupation scale items:* 3, 4, 9, 18, 21, 25.

*Oral Control subscale items:* 2, 5, 8, 13, 15, 19, 20.

Because denial can be a problem on self-report screening instruments, low scores should not be taken to mean that either clinically significant eating disorders symptoms or a formal eating disorder is not present. Collateral information from parents, teammates, and coaches is useful information that can correct for denial, limited self-disclosure, and social desirability. High scores on self-report measures do not necessarily mean the respondent has an eating disorder; however, it does denote concerns regarding body weight, body shape, and eating. However, if you do have a high score, do not panic. It does not necessarily mean that you have a life-threatening condition and that you will have to immediately seek a form of treatment that may be uncomfortable. If you have a score of 20 or higher, this simply means that you should seek the advice of a qualified mental health professional who has experience with treating eating disorders.

In addition to the EAT-26 questions, identification of those at risk for eating disorders is based on information on the individual's body mass index (BMI) and behavioral symptoms reflective of an eating disorder. Following the methodology described for the Eating Disorder Inventory Referral Form (EDI-RF; Garner, 2004) four behavioral questions are included on this version of the EAT-26 aimed at determining the presence of extreme weight-control behaviors as well as providing an estimate of their frequency. These questions assess self-reported binge eating, self-induced vomiting, use of laxatives, and treatment for an eating disorder over the preceding 6 months. Although these content areas could be assessed in the same format as other items, this would not provide the type of frequency data required to evaluate the extent of the problem. Body Mass Index (BMI) is also computed and used to determine if the person is "significantly underweight" compared to age-matched norms. Generally a referral is recommended if a respondent scores "positively" on the EAT-26 items or meets the threshold on one or more of the behavioral criteria.

All self-report measures require open and honest responses in order to provide accurate information. The fact that most people provide honest responses means that the EAT-26 usually provides very useful information about the eating symptoms and concerns that are common in eating disorders.

# Interpreting Eating Attitudes Test (Eat-26)<sup>®</sup> Scores

David M. Garner, Ph.D.  
(Suitable for On-Line Feedback)

The Eating Attitudes Test (EAT-26) is probably the most widely used test used to assess "eating disorder risk" based on attitudes, feelings, and behaviors related to eating and eating disorder symptoms. It was used as a screening instrument in the 1998 National Eating Disorders Screening program and has been used in many other studies to identify individuals with possible eating disorders. However, the EAT-26 does not provide a diagnosis of an eating disorder. A diagnosis can only be provided by a qualified health care professional.

The version of the Eating Attitudes Test (EAT-26) you have just completed has three criteria for determining if you should seek further evaluation of your risk of having an eating disorder. These are:

- 1) Your score on the actual EAT test items;
- 2) Low body weight compared to age-matched norms, and
- 3) Behavioral questions indicating possible eating disorder symptoms or recent significant weight loss.

If you meet one or more of these criteria, you should seek an evaluation by a professional who specializes in the treatment of eating disorders.

## 1) Your Eating Attitudes Test (EAT-26) is: \_\_\_\_

A score at or above 20 on the EAT-26 indicates a high level of concern about dieting, body weight or problematic eating behaviors. Because your score is above 20, you should seek an evaluation by a qualified health professional to determine if your score reflects a problem that warrants clinical attention. However, please keep in mind that high scores do not always reflect over-concern about body weight, body shape, and eating. Screening studies have shown that some people with high scores do not have eating disorders. Regardless of your score, if you are suffering from feelings which are causing you concern or interfering with your daily functioning, you should seek an evaluation from a trained mental health professional.

EAT-26 SCORE	Scoring System for the EAT-26					
	Always	Usually	Often	Sometimes	Rarely	Never
Score for questions 1-25	3	2	1	0	0	0
Score for question # 26	0	0	0	1	2	3

Add the scores for each item together for a total score.

## 2) Your Body Mass Index (BMI) is: \_\_\_\_

If your BMI meets the criterion for "underweight", it is an important risk factor for a serious eating disorder. If your EAT-26 score is 20 or more, then this increases your likelihood of having a serious eating disorder. If your BMI indicates that you are neither "underweight" nor "extremely underweight" compared to age/gender-matched norms then you could still have a serious eating disorder. It just means that it is unlikely that you have anorexia nervosa. If you believe that your body weight is a problem, then it would be good for you to consult with a qualified health professional for further clarification. See the note below for further explanation of BMI.

Age	9	10	11	12	13	14	15	16	17	18	19	20	21+
Female (BMI)	14.0	14.5	14.5	15.0	15.5	16.0	16.5	17.0	17.5	18.0	18.0	18.5	19.0
Male (BMI)	14.0	14.5	15.0	15.0	16.0	16.5	17.0	17.5	18.0	18.5	19.0	19.5	20.0

**3) Behavioral Questions:**

If you scored in the any of the checked boxed (✓), you should seek an evaluation from a trained mental health professional:

<b>In the past 6 months have you:</b>	Never	Once a month or less	2-3 times a month	Once a week	2-6 times a week	Once a day or more
Gone on eating binges where you feel that you may not be able to stop?	<input type="checkbox"/>	<input type="checkbox"/>	✓	✓	✓	✓
Ever made yourself sick (vomited) to control your weight or shape?	<input type="checkbox"/>	✓	✓	✓	✓	✓
Ever used laxatives, diet pills or diuretics (water pills) to control your weight or shape?	<input type="checkbox"/>	✓	✓	✓	✓	✓
Exercised more than 60 minutes a day to lose or to control your weight?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	✓
Lost 20 pounds or more in the past 6 months	Yes	✓	No	<input type="checkbox"/>		

Please remember that the EAT-26 does not provide a diagnosis of an eating disorder. A diagnosis can only be provided by a qualified health care professional.

\* Note on BMI: The EAT-26 includes specific questions on height, weight and gender that can be used to compute Body Mass Index (BMI) for the purpose of determining if you are "at risk" for an eating disorder because your body weight is extremely underweight according to age-matched population norms. BMI is a formula for estimating body mass that takes both height and weight into account. It is calculated by dividing weight (in kilograms) by height in meters, and then divided again by height in meters (kg/m<sup>2</sup>). Alternatively, BMI can be calculated as weight (in pounds) divided by height in inches, then divided again by height in inches and multiplied by 703. We recommend that you seek a professional evaluation for a possible eating disorder if your body weight is "extremely underweight" according to age-matched population norms.

Although BMI is a convenient and useful weight classification tool, it does have limitations. For example, BMI can overestimate fatness for people who are athletic. Also, some races, ethnic groups, and nationalities have different body fat distributions and body compositions; therefore, the norms used are not appropriate for all groups.

**More Information on BMI**

The National Health and Nutrition Examination Survey III (NHANES III, Kuczmarski, Ogden, et al., 2002) has collected reference data to establish weight and height norms at different ages for girls/women and boys/men from birth to 20 years old. These norms indicate that BMI varies considerably with age and gender with children between 5 to 8 years old having the lowest BMI values followed by a steady increase with age. The expected changes in BMI associated females and males as "underweight" (BMI between the 5<sup>th</sup> and 10<sup>th</sup> percentile for girls/women and boys/men from 9 to 20 years old) and "very underweight" (BMI less than the 5<sup>th</sup> percentile). A BMI cutoff of between the 5<sup>th</sup> and 10<sup>th</sup> percentile for different ages and sexes should be used to determine if you meet the "underweight" BMI referral criterion for referral. For men and women 21 years old and older, the "underweight" category according to the NHLBI (1998) survey data were used to determine the "underweight" criterion for referral.

You can easily determine if you meet the BMI thresholds in Table 1 by finding your height on the column on the left in Table 2 and the BMI on the bottom and follow the height and the BMI columns to where the intersect. This is the weight that you need to be at or below for the BMI you have selected.

Although BMI is a convenient and useful weight classification tool, it does have limitations. For example, BMI can overestimate fatness for people who are athletic. Also, some races, ethnic groups, and nationalities have different body fat distributions and body compositions; therefore, the NHANES data are not appropriate for all groups (Kuczmarski, Ogden, et al., 2002).

Age	9	10	11	12	13	14	15	16	17	18	19	20	20+
<b>Female</b>													
Very Underweight (less than or equal to)	13.5	14.0	14.0	14.5	15.0	15.5	16.0	16.5	17.0	17.5	17.5	17.5	18.5
Underweight (between)	13.6-14.0	14.1-14.5	14.1-14.5	14.6-15.0	15.1-15.5	15.6-16.0	16.1-16.5	16.6-17.0	17.1-17.5	17.6-18.0	17.6-18.0	17.6-18.5	18.6-19.0
<b>Male</b>													
Very Underweight (less than or equal to)	13.5	14.0	14.5	14.5	15.0	16.0	16.5	17.0	17.5	18.0	18.5	19.0	19.5
Underweight (between)	13.6-14.0	14.1-14.5	14.6-15.0	14.6-15.0	15.1-16.0	16.1-16.5	16.6-17.0	17.1-17.5	17.6-18.0	18.1-18.5	18.6-19.0	19.1-19.5	19.6-20.0

Data from the NHANES III survey, Kuczmarski, Ogden, et al., 2002.

**Table 2 Body Weight and Height to Calculate Body Mass Index (BMI)**

Height (in.)	Weight (lb.)													
50	50	52	54	55	57	59	60	62	64	66	68	70	78	89
51	52	54	56	58	59	61	63	65	67	68	70	73	81	91
52	54	56	58	60	62	64	65	67	69	71	73	76	85	96
53	56	58	60	62	64	66	68	70	72	74	76	79	88	100
54	58	60	62	64	66	69	71	73	75	77	79	82	91	104
55	60	63	65	67	69	71	73	76	78	80	82	85	95	108
56	63	65	67	69	72	74	76	78	81	83	85	88	98	111
57	65	67	70	72	74	76	79	81	83	86	88	91	101	115
58	67	70	72	74	77	79	82	84	86	89	91	94	105	119
59	70	72	75	77	79	82	84	87	89	92	94	97	108	124
60	72	74	77	80	82	85	87	90	92	95	97	100	112	128
61	74	77	80	82	85	88	90	93	96	98	100	104	116	132
62	77	80	82	85	88	90	93	96	99	101	104	107	120	136
63	79	82	85	88	91	93	96	99	102	105	107	110	124	141
64	82	85	88	91	93	96	99	102	105	108	110	114	128	145
65	84	87	90	93	96	99	102	105	108	112	114	118	132	150
66	87	90	93	96	99	102	106	109	112	115	118	121	136	155
67	90	93	96	99	102	106	109	112	115	118	121	125	140	160
68	92	96	99	102	105	109	112	115	119	122	125	128	145	165
69	95	98	102	105	109	112	115	119	122	126	128	132	148	170
70	98	101	105	108	112	115	119	122	126	129	132	136	153	175
71	101	104	108	111	115	118	122	126	129	133	136	140	157	180
72	103	107	111	114	118	122	125	129	133	137	140	144	162	185
73	106	110	114	118	122	125	129	133	137	140	144	148	166	190
74	109	113	117	121	125	129	133	136	140	144	148	152	171	195
75	112	116	120	124	128	132	136	140	144	148	152	156	175	200
76	115	120	124	128	132	136	140	144	148	152	156	160	180	205
BMI (kg/m)	14.0	14.5	15.0	15.5	16.0	16.5	17.0	17.5	18.0	18.5	19.0	19.5	22.0	25.0

Table 3: 3d, 5 <sup>th</sup> and 10 <sup>th</sup> Percentiles for Females and Males by age from the NHANES						
Age	Female			Male		
	Percentile			Percentile		
	3d	5th	10th	3d	5th	10th
9	13.5	13.7	14.2	13.7	14.0	14.3
10	13.7	14.0	14.5	14.0	14.2	14.6
11	14.1	14.4	14.9	14.3	14.6	15.0
12	14.5	14.8	15.4	14.6	14.9	15.4
13	15.0	15.3	15.9	15.1	15.5	16.0
14	15.4	15.8	16.4	15.7	16.0	16.5
15	15.9	16.3	16.9	16.2	16.6	17.1
16	16.4	16.8	17.4	16.8	17.1	17.7
17	16.8	17.2	17.8	17.3	17.7	18.3
18	17.2	17.6	18.2	17.9	18.2	18.9
19	17.4	17.8	18.4	18.3	18.7	19.4
20	17.4	17.8	18.5	18.7	19.1	19.8

<http://www.cdc.gov/nchs/about/major/nhanes/growthcharts/datafiles.htm>

#### References

- NHLBI (1998). National Heart, Lung and Blood Institute, Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults, June 17, 1998.
- Dotti,A., & Lazzari,R. (1998). Validation and reliability of the Italian EAT-26. *Eating and Weight Disorders*, 3), 188-194.
- Garner, D.M. (1993). Self-report measures for eating disorders. *Current Content, Social and Behavioral Sciences*, 8, 8 Feb. 22 1993, *CC Arts and Humanities*, 5, 20, Mar. 1, 1993.
- Garner, D. M. (2004). *The Eating Disorder Inventory-3 Professional Manual*. Odessa FL: Psychological Assessment Resources Inc.
- Garner, D.M., Rosen, L. and Barry, D. (1998). Eating Disorders in Athletes (839-857). In: *Child and Adolescent Psychiatric Clinics of North America.*, 7, New York: W.B. Saunders.
- Garner, D.M., & Garfinkel, P.E. (1979). The Eating Attitudes Test: an index of the symptoms of anorexia nervosa. *Psychological Medicine*, 9, 273-279.
- Garner, D.M., Olmsted, M.P., Bohr, Y. and Garfinkel, P.E. (1982) The eating attitudes test: Psychometric features and clinical correlates. *Psychological Medicine*, 12, 871-878.
- Kuczmarski, R. J., Ogden, C. L., Guo, S. S., Grummer-Strawn, L. M., Flegal, K. M., Mei, Z., Wei, R., Curtin, L. R., Roche, A. F., & Johnson, C. L. 2000 CDC Growth Charts for the United States: Methods and development. Vital and Health Statistics, Series 11. 246, 1-190. 2002. U.S. National Center for Health Statistics.
- Lee, S., Kwok, K., Liau, C., & Leung, T. (2002). Screening Chinese patients with eating disorders using the Eating Attitudes Test in Hong Kong. *International Journal of Eating Disorders*, 32, 91-97.
- Mintz, L. B., & O'Halloran, M. S. (2000). The Eating Attitudes Test: Validation with DSM-IV eating disorder criteria. *Journal of Personality Assessment*, 74, 489-503.
- Patton, G. C., Johnson-Sabine, E., Wood, K., Mann, A. H., & Wakeling, A. (1990). Abnormal eating attitudes in London schoolgirls: A prospective epidemiological study-outcome at twelve month follow-up. *Psychological Medicine*, 20, 383-394.